

MATH 2352 Solution Sheet 10

BY XIAOYU WEI

Created on May 24, 2015

[Problems] 7.7: 6, 12, 13; 7.8: 7, 18, 22 7.9: 1, 3.

7.7 - 6. For

$$x' = \begin{pmatrix} -1 & 1 \\ -4 & -1 \end{pmatrix} x,$$

(a) Find a fundamental matrix for the given system of equations.

(b) Also find the fundamental matrix $\Phi(t)$ satisfying $\Phi(0) = I$.

Solution.

(a) The general solution is

$$x(t) = c_1 \begin{bmatrix} e^{-t} \cos(2t) \\ -2e^{-t} \sin(2t) \end{bmatrix} + c_2 \begin{bmatrix} e^{-t} \sin(2t) \\ 2e^{-t} \cos(2t) \end{bmatrix}.$$

Then a fundamental matrix can be formed using any two linearly independent solutions, for example,

$$\Psi(t) = \begin{bmatrix} e^{-t} \sin(2t) & e^{-t} \cos(2t) \\ 2e^{-t} \cos(2t) & -2e^{-t} \sin(2t) \end{bmatrix}.$$

(b) The fundamental matrix can be normalized using properly chosen c_1, c_2 ,

$$\Phi(t) = \begin{bmatrix} e^{-t} \cos(2t) & \frac{1}{2} e^{-t} \sin(2t) \\ -2e^{-t} \sin(2t) & e^{-t} \cos(2t) \end{bmatrix},$$

which satisfies $\Phi(0) = I$.

7.7 - 12. Solve the initial value problem

$$x' = \begin{pmatrix} 2 & -1 \\ 3 & -2 \end{pmatrix} x,$$

$$x(0) = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$$

by using the fundamental matrix $\Phi(t)$ for the following system of equations:

$$x' = \begin{pmatrix} 2 & 3 \\ -1 & -2 \end{pmatrix} x.$$

Solution. We intend to use the relation $x(t) = \Phi(t)x(0)$.

To solve for $\Phi(t)$, we directly compute

$$\Phi(t) = e^{At},$$

where

$$A = \begin{pmatrix} 2 & -1 \\ 3 & -2 \end{pmatrix}.$$

A can be decomposed as Jordan canonical form

$$\begin{aligned} A &= SJS^{-1} \\ &= \begin{pmatrix} -1 & -3 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{2} & \frac{3}{2} \\ -\frac{1}{2} & -\frac{1}{2} \end{pmatrix}, \end{aligned}$$

which can be obtained from the transposed matrix.

Then

$$\begin{aligned} e^{At} &= Se^{Jt}S^{-1} \\ &= \begin{pmatrix} -1 & -3 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} e^{-t} & 0 \\ 0 & e^t \end{pmatrix} \begin{pmatrix} \frac{1}{2} & \frac{3}{2} \\ -\frac{1}{2} & -\frac{1}{2} \end{pmatrix}. \end{aligned}$$

To compute $x(t) = \Phi(t)x(0)$, we have

$$\begin{aligned} x(t) &= \begin{pmatrix} -1 & -3 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} e^{-t} & 0 \\ 0 & e^t \end{pmatrix} \begin{pmatrix} \frac{1}{2} & \frac{3}{2} \\ -\frac{1}{2} & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix} \\ &= \begin{pmatrix} -1 & -3 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} e^{-t} & 0 \\ 0 & e^t \end{pmatrix} \begin{pmatrix} \frac{11}{2} \\ -\frac{5}{2} \end{pmatrix} \\ &= \begin{pmatrix} -1 & -3 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \frac{11}{2}e^{-t} \\ -\frac{5}{2}e^t \end{pmatrix} \\ &= \begin{pmatrix} -\frac{11}{2}e^{-t} + \frac{15}{2}e^t \\ \frac{11}{2}e^{-t} - \frac{5}{2}e^t \end{pmatrix}. \end{aligned}$$

7.7 - 13. Show that $\Phi(t) = \Psi(t)\Psi^{-1}(t_0)$, where $\Phi(t)$ and $\Psi(t)$ are as defined in this section (or as in the slides).

Solution. It is equivalent to show

$$\Phi(t)\Psi(t_0) = \Psi(t).$$

Consider each column of the equation above, it means that the solution forming the fundamental matrix is linear combination of columns of $\Phi(t)$, with coefficients as initial values. This is exactly the definition of $\Phi(t)$.

7.8 - 7. Consider the initial value problem

$$x' = \begin{pmatrix} 1 & -4 \\ 4 & -7 \end{pmatrix} x,$$

$$x(0) = \begin{pmatrix} 4 \\ 2 \end{pmatrix}.$$

(a) Find the solution.

(b) Draw the trajectory of the solution in the x_1x_2 -plane, and also draw the graph of x_1 versus t .

Solution.

(a) Same as the previous problem,

$$\begin{aligned} x(t) &= \Phi(t)x(0) \\ &= e^{At}x(0) \quad (\text{Jordan canonical form}) \\ &= Se^{Jt}S^{-1}x(0) \\ &= \begin{pmatrix} 1 & \frac{1}{4} \\ 1 & 0 \end{pmatrix} \begin{pmatrix} e^{-3t} & te^t \\ 0 & e^{-3t} \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 4 & -4 \end{pmatrix} \begin{pmatrix} 4 \\ 2 \end{pmatrix} \\ &= \begin{pmatrix} 1 & \frac{1}{4} \\ 1 & 0 \end{pmatrix} \begin{pmatrix} e^{-3t} & te^t \\ 0 & e^{-3t} \end{pmatrix} \begin{pmatrix} 2 \\ 8 \end{pmatrix} \\ &= \begin{pmatrix} 1 & \frac{1}{4} \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 2e^{-3t} + 8te^t \\ 8e^{-3t} \end{pmatrix} \\ &= \begin{pmatrix} 4e^{-3t} + 8te^t \\ 2e^{-3t} + 8te^t \end{pmatrix}. \end{aligned}$$

(b) As shown below:




Figure 1. Trajectory (left) and $x_1(t)$ (right)

7.8 - 18. Consider the system

$$x' = Ax = \begin{pmatrix} 1 & 1 & 1 \\ 2 & 1 & -1 \\ -3 & 2 & 4 \end{pmatrix} x. \quad (1)$$

(a) Show that $r=2$ is an eigenvalue of algebraic multiplicity 3 of the coefficient matrix A and that there is only one corresponding eigenvector, namely,

$$\xi^{(1)} = \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}.$$

(b) Using the information in part (a), write down one solution $x^{(1)}(t)$ of the system (1). There is no other solution of the purely exponential form $x = \xi e^{rt}$.

(c) To find a second solution, assume that $x = \xi t e^{2t} + \eta e^{2t}$. Show that ξ and η satisfy the equations

$$\begin{aligned} (A - 2I) \xi &= 0, \\ (A - 2I) \eta &= \xi. \end{aligned}$$

Since ξ has already been found in part (a), solve the second equation for η . Neglect the multiple of $\xi^{(1)}$ that appears in η , since it leads only to a multiple of the first solution $x^{(1)}$. Then write down a second solution $x^{(2)}(t)$ of the system (1).

(d) To find a third solution, assume that $x = \xi (t^2/2) e^{2t} + \eta t e^{2t} + \zeta e^{2t}$. Show that ξ , η and ζ satisfy the equations

$$\begin{aligned} (A - 2I) \xi &= 0, \\ (A - 2I) \eta &= \xi, \\ (A - 2I) \zeta &= \eta. \end{aligned}$$

The first two equations are the same as in part (c), so solve the third equation for ζ , again neglecting the multiple of $\xi^{(1)}$ that appears. Then write down a third solution $x^{(3)}(t)$ of the system (1).

(e) Write down a fundamental matrix $\Psi(t)$ for the system (1).

(f) Form a matrix T with the eigenvector $\xi^{(1)}$ in the first column and the generalized eigenvector η and ζ in the second and third columns. Then find T^{-1} and form the product $J = T^{-1}AT$. The matrix J is the Jordan form of A .

Solution.

The solution procedure is to evaluate the Jordan canonical form and solve the ODE in parallel. Following the instructions, we can get

$$A = \begin{pmatrix} 0 & -1 & -2 \\ -1 & -1 & -3 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} 0 & 0 & 1 \\ -3 & 2 & 2 \\ -1 & -1 & -1 \end{pmatrix},$$

and

$$x(t) = c_1 \begin{pmatrix} -2e^{2t}(t-1) \\ -e^{2t}(t-4)t \\ e^{2t}(t-6)t \end{pmatrix} + c_2 \begin{pmatrix} 2e^{2t}t \\ e^{2t}[(t-2)t+2] \\ -e^{2t}(t-4)t \end{pmatrix} + c_3 \begin{pmatrix} 2e^{2t}t \\ e^{2t}(t-2)t \\ -e^{2t}[(t-4)t-2] \end{pmatrix}.$$

7.8 - 22. Let

$$J = \begin{pmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 1 \\ 0 & 0 & \lambda \end{pmatrix},$$

where λ is an arbitrary real number.

(a) Find J^2 , J^3 , and J^4 .

(b) Use an inductive argument to show that

$$J^n = \begin{pmatrix} \lambda^n & n\lambda^{n-1} & [n(n-1)/2]\lambda^{n-2} \\ 0 & \lambda^n & n\lambda^{n-1} \\ 0 & 0 & \lambda^n \end{pmatrix}.$$

(c) Determine e^{Jt} .

(d) Note that if you choose $\lambda = 2$, then the matrix J in this problem is the same as the matrix J in Problem 18(f), form the product $T e^{Jt}$ with $\lambda = 2$. The resulting matrix is the same as the fundamental matrix $\Psi(t)$ in Problem 18(e).

Solution. The solution is straight forward following the instructions, and to determine e^{Jt} , consider its (formal) Taylor expansion.

$$e^{Jt} = I + Jt + \frac{1}{2}J^2 t^2 + \dots$$

7.9 - 1. Find the general solution of the given system of equations.

$$x' = \begin{pmatrix} 2 & 3 \\ -1 & -2 \end{pmatrix} x + \begin{pmatrix} e^t \\ t \end{pmatrix}.$$

Solution. Solve the homogeneous equation first, then find a special solution to the inhomogeneous equation.

See the solution to the next problem for step-by-step guidelines.

7.9 - 3. Find the general solution of the given system of equations.

$$x' = \begin{pmatrix} 2 & 1 \\ -5 & -2 \end{pmatrix} x + \begin{pmatrix} -\cos t \\ \sin t \end{pmatrix}.$$

Solution.

1. Find the eigenvalues and eigenvectors of the matrix

$$A = \begin{pmatrix} 2 & 1 \\ -5 & -2 \end{pmatrix}.$$

We have

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i, \end{aligned}$$

and

$$v_1 = \begin{pmatrix} -\frac{2}{5} - \frac{i}{5} \\ 1 \end{pmatrix},$$

$$v_2 = \begin{pmatrix} -\frac{2}{5} + \frac{i}{5} \\ 1 \end{pmatrix},$$

2. Compute the solution to the homogeneous equation

$$\begin{aligned}x_1 &= e^{\lambda_1 t} v_1 \\&= \begin{pmatrix} -\frac{2}{5} - \frac{i}{5} \\ 1 \end{pmatrix} e^{it} \\x_2 &= e^{\lambda_2 t} v_2 \\&= \begin{pmatrix} -\frac{2}{5} + \frac{i}{5} \\ 1 \end{pmatrix} e^{-it}.\end{aligned}$$

And let $X(t) = [\operatorname{Re}(x_1), \operatorname{Im}(x_1)]$ be a (real) fundamental matrix.

$$X(t) = \begin{pmatrix} -\frac{2}{5} \cos t + \frac{1}{5} \sin t & -\frac{2}{5} \sin t - \frac{1}{5} \cos t \\ \cos t & \sin t \end{pmatrix}$$

3. Find a particular real solution to the nonhomogeneous equation by

$$x^*(t) = X(t) \int X^{-1}(t) g(t) dt,$$

where g is the source term

$$g(t) = \begin{pmatrix} -\cos t \\ \sin t \end{pmatrix}.$$

4. The general solution is

$$x(t) = x^*(t) + X(t) C,$$

where C is a constant vector.